

## Supplementary Files

Supplementary Table A Clinical presentations

Clinical presentations	n (%)
Lymphadenopathy	31 (52)
Abdominal lump	14 (23)
Intestinal obstruction	3 (5)
Testicular swelling	2 (3)
Swelling jaw	2 (3)
Leg swelling	2 (3)
Leptomeningeal disease	1(2)
Raised Intracranial tension	1(2)
Breast lump	1(2)
Scalp swelling	1(2)
Nasopharyngeal swelling	1(2)
Chest wall swelling	1(2)
Vaginal mass	1(2)

Supplementary Table B Hemogram of 11 patients with Burkitt's leukemia

No.	HB	TC	PLC
1	13.5	13,000	4,09,000
2	9.7	12,500	38,000
3	9.4	6,300	2,61,000
4	10.2	10,500	48,000
5	7.3	10,000	1,0,5000
6	8.3	4,800	40,000
7	8.2	29,400	2,31,000
8	10.6	30,200	2,35,000
9	10.2	7,800	37,000
10	8.2	2,300	33,000
11	14.6	16,000	1,23,000

Abbreviations: HB, hemoglobin; PLC, platelet count; TLC, total WBC count.

Supplementary Table C Univariate Cox regression analysis

Variables	PFS				OS			
	p-Value	HR	95.0% CI for HR		p-Value	HR	95.0% CI for HR	
			Lower	Upper			Lower	Upper
Age group 40–65 vs. <40	0.012	3.72	1.33	10.35	0.01	3.68	1.324	10.23
>65 vs. <40	0.001	10.31	2.91	36.53	0.00	10.1	2.87	35.95
Sex (female vs. male)	0.314	0.603	0.22	1.616	0.31	0.60	0.224	1.614
LDH ( $\geq$ 2,000 vs. <2,000)	0.58	0.732	0.24	2.207	0.56	0.72	0.24	2.182
Burkitt's leukemia vs. Burkitt's lymphoma	0.887	1.074	0.40	2.879	0.87	1.08	0.403	2.895
Stage (2 vs. 1)	0.464	1.461	0.52	4.035	0.46	1.45	0.528	4.024
Stage (3 vs. 1)	0.981	0	0	2.597	0.98	0	0	2.584
Stage (4 vs. 1)	0.947	0.967	0.36		0.93	0.96	0.358	
Hyper-CVAD $\pm$ R vs. CHOP/CVP $\pm$ R	0.139	0.519	0.21	1.238	0.14	0.52	0.218	1.242

Abbreviations: CI, confidence interval; HR, hazard ratio; PFS, progression-free survival; OS, overall survival.